

Sources

US Census 2007.

National Health Interview Survey, 2006.

Lusk, R., et al, Laryngoscope 2006; 116: 2099-2107.

Benninger, M., Otolaryngol Head Neck Surg 2003; 129S: S1-S32.

Gliklich, R., et al, J Allergy Clin Immunol 1999; 113: 104-109.

Ray, N., et al, J Allergy Clin Immunol 1999; 103: 408-414.

Collins, J., Vital Health Stat 1997; 10: 1-89.

Hamilos, J Allergy Clin Immunol 2000; 106: 213-227.

Stankiewicz, J., et al, Am J Rhinol 2003; 17(3): 139-142.

Subramanian, H., et al, Am J Rhinol 2002; 16(6): 303-312.

Hessler, J., et al, Am J Rhinol 2007; 21(1): 10-18.

Levine, H., et al, Annals Oto Rhino Laryngol 2008; 117(4): 263-270.

Weiss, R., et al, Otolaryngol Head Neck Surg 2008; 139: S38-S46.

Bolger W., et al, Otolaryngology Head and Neck Surgery, 2007; 137: 10-20.

Kuhn F.A., et al, Otolaryngology Head and Neck Surgery, 2008; 139: 3S3, S27-S37.

Friedman M., Schalch P, et al. American Journal of Rhinology, 2008; 22: No.2.

Ramadan H.H., Annals of Otology, Rhinology and Laryngology, 2009; 118(3):161-165.

Church C., et al, Otolaryngology Head and Neck Surgery, 2008; 138: 187-191.